

JOURNAL OF ALGEBRA **88**, 63–67 (1984)

Bireflectionality of the Weak Orthogonal and the Weak Symplectic Groups

E. W. ELLERS*

*Department of Mathematics, University of Toronto,
Toronto, Ontario M5S 1A1, Canada*

AND

R. FRANK AND W. NOLTE

*Fachbereich Mathematik, AG 2 Geometrie und Algebra, Technische Hochschule Darmstadt,
Schlossgartenstrasse 7, D-6100 Darmstadt, West Germany*

Communicated by B. Huppert

Received October 18, 1982

1. INTRODUCTION

Let V be a finite-dimensional vector space, Q a quadratic form and $f_Q = f$ the bilinear form associated with Q . We assume (V, f_Q) is regular. Then the orthogonal group $O(V)$ is bireflectional, i.e., every isometry in $O(V)$ is a product of two involutory isometries in $O(V)$. This has been shown in [8] if the field of scalars K has characteristic distinct from 2 and in [4] and [5] if $\text{char } K = 2$. The latter papers also establish the bireflectionality for the symplectic group $\text{Sp}(V)$, again under the assumption that (V, f) is regular and $\text{char } K = 2$. For $\text{char } K \neq 2$ the symplectic group is not bireflectional (see [3]).

We shall extend the results just mentioned. We shall drop the assumptions that V is finite dimensional and that (V, f) is regular, i.e., the vector space V may be infinite dimensional and the radical of V may be distinct from zero.

We shall use the notation and the concepts in [2]. For every $\pi \in \text{Hom}(V, V)$ we define $F(\pi) = \{v \in V; v\pi = v\}$ and $B(\pi) = \{v\pi - v; v \in V\}$. The spaces $F(\pi)$ and $B(\pi)$ are called fix and path of π , respectively.

The groups $O^*(V) = \{\pi \in O(V); \text{rad } V \subset F(\pi) \text{ and } \dim B(\pi) < \infty\}$ and $\text{Sp}^*(V) = \{\pi \in \text{Sp}(V); \text{rad } V \subset F(\pi) \text{ and } \dim B(\pi) < \infty\}$ are called the weak orthogonal and the weak symplectic group, respectively.

* Research supported in part by NSERC Canada Grant No. A7251.

In Section 2 we shall assume that the dimension of the vector space V is finite. We shall show that under this assumption the weak orthogonal group and for $\text{char } K = 2$ also the weak symplectic group are bireflectional. Since for $\text{char } K \neq 2$ the symplectic group is not bireflectional, we shall always assume that $\text{char } K = 2$ when we deal with the weak symplectic group $\text{Sp}^*(V)$.

In Section 3 we investigate the situation in infinite-dimensional vector spaces. First we see in Lemma 3 that for $\text{char } K \neq 2$ every involution in $O^*(V)$ and therefore every product of two involutions is a product of reflections. Since there are elements in $O^*(V)$ which are not products of reflections (see [2, p. 116]) we see that $O^*(V)$ is not bireflectional. This suggests to define $O^{**}(V)$ as the subgroup of $O(V)$ whose elements are products of reflections and to give an analogous definition for $\text{Sp}^{**}(V)$. We shall show in Theorem 4 that $O^{**}(V)$ and $\text{Sp}^{**}(V)$ are bireflectional if $K \neq GF(2)$.

Even if $K = GF(2)$, we see in Theorem 4 that the elements in $O^{**}(V)$ and $\text{Sp}^{**}(V)$ are products of two involutions in $O^*(V)$ and $\text{Sp}^*(V)$, respectively.

Finally we need a few formulas whose proof is not difficult (see, e.g., [7]):

For $\pi_1, \pi_2 \in \text{Hom}(V, V)$ and $\pi \in GL(V)$ we get

$$B(\pi_1 \pi_2) \subset B(\pi_1) + B(\pi_2), \quad (1)$$

$$B(\pi^{-1}) = B(\pi), \quad (2)$$

$$B(\pi^m) \subset B(\pi) \quad \text{for all integers } m. \quad (3)$$

2. THE GROUPS $O^*(V)$ AND $\text{Sp}^*(V)$ FOR A FINITE-DIMENSIONAL VECTOR SPACE V

In this section we assume that V is a finite-dimensional vector space. For the proof of Theorem 1 we embed V into a regular vector space \bar{V} . Then we apply results of [4] and [8]. Before we can do that it is necessary to learn more about the involutory factors ρ_1 and ρ_2 of π which were constructed in [4] and [8]. We need the following result:

Assume V is regular. Then every isometry π is a product of two involutory isometries ρ_1 and ρ_2 such that

$$B(\rho_i) \subset B(\pi), \quad i = 1, 2. \quad (4)$$

We shall see that (4) is true for the involutory isometries ρ_1 and ρ_2 in [4] (see Lemma 1 and the proof of Lemma 4) and [8] (see the proofs of Lemmas 2 and 5). The vector space V is an orthogonal sum of regular π -cyclic and regular π -bicyclic subspaces. Since a π -bicyclic subspace is a direct sum of two (nonregular) π -cyclic subspaces, we get that V is a direct

sum of π -cyclic subspaces $W_k: V = \sum_{k=1}^q W_k$ for some integer q . Let w_k be a π -cyclic generator of W_k and assume $\dim W_k = n_k$. Then $\{w_k \pi^i; k = 1, \dots, q; i = 0, \dots, n_{k-1}\}$ is a basis for V .

First we shall deal with [4]. The restriction of ρ_1 to the π -cyclic subspace W_k is one of the two involutory mappings given in [4], (7), which we now call σ_1 and σ_2 . We give their definitions (putting $w_k = w$ and $n_k = n$):

$$w\pi^i\sigma_1 = w\pi^{n-i} \quad \text{and} \quad w\pi^i\sigma_2 = w\pi^{n+1-i} \quad \text{for all integers } i. \quad (5)$$

Then $w(\pi^i\sigma_1 - \pi^i) = w(\pi^{n-i} - \pi^i) = w\pi^i(\pi^{n-2i} - 1) \in B(\pi^{n-2i}) \subset B(\pi)$ by (3) and consequently $B(\sigma_1) \subset B(\pi)$. A similar calculation shows $B(\sigma_2) \subset B(\pi)$. Therefore $B(\rho_1) \subset B(\pi)$.

Second we consider [8] (i.e., $\text{char } K \neq 2$). In [8] the path of ρ_1 is always generated by vectors of the form $u(\pi^i - \pi^{-i})$ for some $u \in V$. Now $u(\pi^i - \pi^{-i}) = u\pi^{-i}(\pi^{2i} - 1) \in B(\pi^{2i}) \subset B(\pi)$. Thus again $B(\rho_1) \subset B(\pi)$.

Using (2) we get $B(\rho_2) \subset B(\rho_1) + B(\pi) \subset B(\pi)$ and therefore (4) holds.

Now we are ready to state and prove our first result. We allow V to be nonregular.

THEOREM 1. *Assume V is a finite-dimensional vector space. Then the weak orthogonal group $O^*(V)$ is bireflectional and if $\text{char } K = 2$, also the weak symplectic group $\text{Sp}^*(V)$ is bireflectional.*

Proof. Let $V = A \oplus \text{rad } V$ (then A is regular) and $k = \dim \text{rad } V$. Then V can be embedded into a regular vector space \bar{V} such that $\dim \bar{V} = \dim V + k$. By Witt's theorem (see [1, p. 71]) every isometry π of V can be extended to an isometry $\bar{\pi}$ of \bar{V} . In \bar{V} we have $k + \dim(\text{rad } V)^\perp = \dim \text{rad } V + \dim(\text{rad } V)^\perp = \dim \bar{V} = \dim V + k$. Hence $\dim(\text{rad } V)^\perp = \dim V$. This implies $V = (\text{rad } V)^\perp$ since $V \subset (\text{rad } V)^\perp$. Clearly $\text{rad } V \subset F(\pi) \subset F(\bar{\pi})$ and therefore $V = (\text{rad } V)^\perp \supset F(\bar{\pi})^\perp = B(\bar{\pi})$.

By [4] and [8] there are involutions $\bar{\rho}_1, \bar{\rho}_2 \in O(\bar{V})$ and $\bar{\rho}_1, \bar{\rho}_2 \in \text{Sp}(\bar{V})$, respectively, such that $\bar{\pi} = \bar{\rho}_1 \bar{\rho}_2$. By (4) we have $B(\bar{\rho}_i) \subset B(\bar{\pi}) \subset V$; thus V is invariant under $\bar{\rho}_i$ for $i = 1, 2$. Let $\rho_i = \nu | \bar{\rho}_i$, then $\pi = \rho_1 \rho_2$.

Since \bar{V} is regular and since $B(\bar{\rho}_i) \subset V$ we get $F(\bar{\rho}_i) = B(\bar{\rho}_i)^\perp \supset V^\perp$. Finally $\text{rad } V = V \cap V^\perp \subset V \cap F(\bar{\rho}_i) = F(\rho_i)$.

COROLLARY 2. *Assume V is a finite-dimensional vector space. Then $O^*(V) \cup (-1) \cdot O^*(V)$ is a bireflectional group.*

Proof. Clearly $O^*(V) \cup (-1) \cdot O^*(V)$ is a group. If $\pi \in (-1) \cdot O^*(V)$, then $-\pi \in O^*(V)$. Thus $-\pi = \rho_1 \rho_2$, where $\rho_i \in O^*(V)$ are involutions. Therefore $\pi = (-\rho_1) \rho_2$, $(-\rho_1) \in (-1) \cdot O^*(V)$ and $-\rho_1$ is an involution.

3. THE SUBGROUPS GENERATED BY REFLECTIONS OR SYMPLECTIC TRANSVECTIONS

In this section the dimension of V is arbitrary, it may be infinite.

First we observe that $O^*(V)$ is not bireflectional if $\dim V/R$ is infinite, $R \neq \{0\}$, and $\text{char } K \neq 2$. This is a consequence of the following Lemma 3 and [2, p. 116].

LEMMA 3. *Assume $\text{char } K \neq 2$. If $\rho \in O^*(V)$ is an involution, then ρ is a product of reflections.*

Proof. Since ρ is an involution and since $\text{char } K \neq 2$, we have $B(\rho) \cap F(\rho) = \{0\}$. Hence $B(\rho) \cap \text{rad } V = \{0\}$ since $\text{rad } V \subset F(\rho)$. Therefore $B(\rho)^\perp = F(\rho)$ by [2, p. 105]. Thus $\text{rad } B(\rho) = B(\rho)^\perp \cap B(\rho) = F(\rho) \cap B(\rho) = \{0\}$. Hence $B(\rho)$ is regular. Consequently there is an orthogonal basis $\{a_1, \dots, a_m\}$ for $B(\rho)$ and $\rho = \sigma_1 \cdots \sigma_m$, where σ_i is a reflection such that $B(\sigma_i) = Ka_i$ for $i = 1, \dots, m$.

Lemma 3 is not true for $\text{char } K = 2$. A counterexample can easily be constructed using the example given in [2, p. 116].

Our aim is to show that suitable subgroups of $O^*(V)$ and $\text{Sp}^*(V)$ are bireflectional, namely, the groups $O^{**}(V)$ and $\text{Sp}^{**}(V)$: $O^{**}(V)$ consists of all elements in $O(V)$ that are products of reflections; $\text{Sp}^{**}(V)$ consists of all elements in $\text{Sp}(V)$ that are products of symplectic transvections τ_i such that $B(\tau_i) \cap \text{rad } V = \{0\}$ and $\text{char } K = 2$.

Clearly, $O^{**}(V) \subset O^*(V)$ and $\text{Sp}^{**}(V) \subset \text{Sp}^*(V)$.

THEOREM 4. *For $\pi \in O^{**}(V)$ and $\pi \in \text{Sp}^{**}(V)$ there are involutory isometries $\rho_i \in O^*(V)$ and $\rho_i \in \text{Sp}^*(V)$, respectively, $i = 1, 2$, such that $\pi = \rho_1 \rho_2$. If $K \neq GF(2)$, then $O^{**}(V)$ and $\text{Sp}^{**}(V)$ are bireflectional.*

Proof. By [2, p. 105], there is some subspace $D \subset V$ such that $B(\pi)^\perp = F(\pi) \oplus D$ and $D(\pi - 1) = B(\pi) \cap \text{rad } V$. If there is some $d_1 \in D \setminus \{0\}$, then there is some hyperplane H_1 such that $F(\pi) \subset H_1$ and $d_1 \notin H_1$. Put $D_1 = D \cap H_1$. Then $\dim D_1 = \dim D - 1$ and $D_1 \cap F(\pi) = \{0\}$. Continuing this process we get hyperplanes H_1, \dots, H_m where $m = \dim D$ such that $D \cap H_1 \cap \dots \cap H_m = \{0\}$ and $F(\pi) \subset H_1 \cap \dots \cap H_m$. By [2, Lemma 18], there are $t_i \in V$ such that $H_i = t_i^\perp$ for $i = 1, \dots, m$. Put $T = \langle t_1, \dots, t_m \rangle$. Then $F(\pi) \subset T^\perp$ and $D \cap T^\perp = \{0\}$.

Suppose C is a complement of $F(\pi)$, i.e., $V = F(\pi) \oplus C$. Let $X = B(\pi) + T + C$. Then X is π -invariant, finite dimensional, and $X^\perp = B(\pi)^\perp \cap T^\perp \cap C^\perp = (D + F(\pi)) \cap T^\perp \cap C^\perp = F(\pi) \cap C^\perp \subset F(\pi)$.

There is some finite-dimensional subspace W of V such that $X \subset W$ and $\text{rad } W \subset \text{rad } V$. In order to see that, let $X = Y \oplus \text{rad } X$ and if there is some $x \in \text{rad } X \setminus \text{rad } V$, then $V = Y \oplus Y^\perp$ and there is some $z \in Y^\perp$ such

that $f(x, z) \neq 0$ and therefore $\dim \operatorname{rad}(X + Kz) \leq \dim \operatorname{rad} X - 1$. If $\operatorname{rad}(X + Kz) \subset \operatorname{rad} V$, put $X + Kz = W$. Otherwise repeat the construction.

Now $V = W + W^\perp$ and $W^\perp \subset X^\perp \subset F(\pi)$.

Since $\operatorname{rad} W \subset \operatorname{rad} V$, we have $_{\operatorname{rad} W} \pi = 1$. Using Theorem 1 we get $_{\operatorname{rad} W} \pi = \tilde{\rho}_1 \tilde{\rho}_2$, where for $i = 1, 2$ $\tilde{\rho}_i$ are involutory isometries of W and $_{\operatorname{rad} W} \tilde{\rho}_i = 1$. Therefore we can define the desired involutory isometries ρ_i by $_{\operatorname{rad} W} \rho_i = \tilde{\rho}_i$ and $_{W^\perp} \rho_i = 1$. Clearly, $B(\rho_i) = B(\tilde{\rho}_i)$, $\dim B(\rho_i) < \infty$, $V^\perp \subset W^\perp \subset F(\rho_i)$, so $\rho_i \in O^*(V)$, or $\operatorname{Sp}^*(V)$, and $\rho_1 \rho_2 = \pi$.

Now we assume $K \neq GF(2)$. This together with the fact that $\tilde{\rho}_i \in O^*(W)$ or $\operatorname{Sp}^*(W)$ for the finite-dimensional space W implies by [2, Theorem 33 and 24], respectively, that $\tilde{\rho}_i$ is a product of reflections or regular symplectic transvections, respectively. These reflections and transvections can be extended in a natural way to reflections and transvections of V , respectively. Then ρ_i is a product of these reflections or transvections, respectively. This proves our last contention.

REFERENCES

1. N. BOURBAKI, "Éléments de mathématique," Livre II, Chapitre 9, Formes sesquilineaires et formes quadratiques, Herman, Paris, 1959.
2. E. W. ELLERS, Decomposition of orthogonal, symplectic, and unitary isometries into simple isometries, *Abh. Math. Sem. Univ. Hamburg* **46** (1977), 97–127.
3. E. W. ELLERS, Bireflectionality in classical groups, *Canad. J. Math.* **29** (1977), 1157–1162.
4. E. W. ELLERS AND W. NOLTE, Bireflectionality of orthogonal and symplectic groups, *Arch. Math.* **39** (1982), 113–118.
5. R. GOW, Products of two involutions in classical groups of characteristic 2, *J. Algebra* **71** (1981), 583–591.
6. B. HUPPERT, Isometrien von Vektorräumen I, *Arch. Math.* **35** (1980), 164–176; II, *Math. Z.* **175** (1980), 5–20.
7. O. T. O'MEARA, Group-theoretic characterization of transvections using CDC, *Math. Z.* **110** (1969), 385–394.
8. M. J. WONENBURGER, Transformations which are products of two involutions, *J. Math. Mech.* **16** (1966), 327–338.